

## BIRTH TO AGE 2

### Chickenpox (varicella)

- Between 12 and 15 months

### Diphtheria, tetanus and pertussis (DTaP) *four times*

- 2 months
- 4 months
- 6 months
- between 15 and 18 months

### Flu *Yearly by the end of October, if possible*

- Beginning at 6 months

### Haemophilus influenza (Hib)

*Three or four times depending on brand (common upper respiratory infection that can cause meningitis)*

- 2 months
- 4 months
- 6 months (depending on brand)
- Between 12 and 15 months

### Hepatitis A *Two times*

- Between 12 and 23 months
- Second dose 6 months after the first one

### Hepatitis B *Three times*

- Shortly after birth
- Between 6 and 18 months
- Between 1 to 2 months

### Measles, mumps, rubella (MMR)

- Between 12 and 15 months

### Pneumococcal (PCV13) *Four times*

- 2 months
- 4 months
- 6 months
- Between 12 and 15 months

### Polio (IPV) *Three times*

- 2 months
- 4 months
- Between 6 and 18 months

### Rotavirus (RV)

*Two or three times depending on brand*

#### Rotarix brand

- 2 months
- 4 months

#### RotaTeq brand

- 2 months
- 4 months
- 6 months

### SARS-COV2 (COVID-19): mRNA vaccine

#### Moderna

*Two to three times depending on immune system*

- 6 months to 5 years
- Second dose 4 to 8 weeks after the first one
- Third dose ONLY for moderately or severely immunocompromised populations, 4 weeks after

#### Pfizer *Three times*

- 6 months to 4 years
- Second dose 3 to 8 weeks after the first one
- Third dose at least 8 weeks after the second one

## AGES 3 THROUGH 11

### Chickenpox (varicella)

- Between 4 and 6 years

### Diphtheria, tetanus and pertussis (DTaP)

- Between 4 and 6 years

### Flu *Yearly by the end of October, if possible*

- Every year

### Measles, mumps, rubella (MMR)

- Between 4 and 6 years

### Polio (IPV) *Three times*

- Between 4 and 6 years

### Human papillomavirus (HPV) *two times*

- Between 11 and 12 years
- 6–12 months after first dose

### Meningococcal conjugate *two times*

- Between 11 and 12 years
- Booster at 16 years

### Tetanus, diphtheria and pertussis (Tdap)

*a booster to the Tdap*

- Between 11 and 12 years

### SARS-COV2 (COVID-19): mRNA vaccine

#### Moderna

*Two to three times depending on immune system*

- 6 to 11 years
- Second dose 4 to 8 weeks after the first one
- Third dose ONLY for moderately or severely immunocompromised populations, 4 weeks after second dose

#### Pfizer *Three to four times depending on immune system*

- 5 to 11 years
- Second dose 3 to 8 weeks after the first one
- Third dose at least 8 weeks after the second one
- Fourth dose ONLY for moderately or severely immunocompromised populations, 3 months after third dose

## AGES 12 THROUGH 18

### Flu *Yearly by the end of October, if possible*

- Every year

### Meningococcal serogroup B

- Between 16 and 23 years

### SARS-COV2 (COVID-19): mRNA vaccine

#### Moderna

*Two to three times depending on immune system*

- 12 to 17 years
- Second dose 4 to 8 weeks after the first one
- Third dose ONLY for moderately or severely immunocompromised populations, 4 weeks after second dose

#### Pfizer *Three to four times depending on immune system*

- 12 to 17 years
- Second dose 3 to 8 weeks after the first one
- Third dose at least 8 weeks after the second one
- Fourth dose ONLY for moderately or severely immunocompromised populations, 3 months after third dose